

EUCRISA: NOW APPROVED FOR USE IN PATIENTS AS YOUNG AS 3 MONTHS OF AGE WITH MILD-TO-MODERATE ATOPIC DERMATITIS (AD)¹

eucrisa
crisaborole ointment 2%



EUCRISA IS THE FIRST AND ONLY FDA-APPROVED NONSTEROIDAL TOPICAL Rx MEDICATION FOR AD IN PATIENTS AGED 3 MONTHS AND OLDER AND CAN BE USED AS PART OF A LONG-TERM TREATMENT PLAN^{1-3*†‡}

In pivotal trials of patients 2 years and older

Significantly more EUCRISA patients achieved success in ISGA[§] at Day 29 in two clinical trials^{1,4,5}

EUCRISA (n=503) 32.8%, vehicle (n=256) 25.4%; P=0.038 in Trial 1

EUCRISA (n=513) 31.4%, vehicle (n=250) 18.0%; P<0.001 in Trial 2

The most common adverse reaction occurring in ≥1% of patients in pivotal trials was application site pain[¶]

Occurred in 4% (45/1012) of patients treated with EUCRISA vs 1% (6/499) for vehicle[¶]

—Application site pain resolved within 1 day for 77.6% of patients who reported it[¶]

Discontinuation rates due to adverse events were 1.2% for both EUCRISA and vehicle in a pooled analysis[¶]

Study Design^{1,4}

Two multicenter, randomized, double-blind, vehicle-controlled trials (Trial 1 and Trial 2) treating 1522 patients (1016 EUCRISA; 506 vehicle), 2 to 79 years of age, with mild-to-moderate AD. Patients were instructed to apply EUCRISA or vehicle twice daily for 28 days. Efficacy and safety endpoints were evaluated at Days 1 (baseline), 8, 15, 22, and 29. The primary efficacy endpoint was success in ISGA at Day 29.

*Should be applied twice daily to the affected areas.¹

[†]Patients aged 2 years and older were studied in pivotal trials for 28 days and in an open-label safety study for up to 48 weeks. In the open-label safety extension study, 517 patients aged 2 and older were cycled on or off therapy in 4-week intervals as needed for 48 weeks.³

[‡]'Long-term treatment plan' refers to your general skin care recommendations for patients with AD, including treatment recommendations for the recurrence of signs and symptoms of AD on affected skin.

[§]Success in ISGA, a stringent metric, is defined as Clear (0) or Almost Clear (1) AND at least a 2-grade improvement from baseline at Day 29.¹

[¶]Refers to skin sensations such as burning or stinging.

ISGA=Investigator's Static Global Assessment.

IMPORTANT SAFETY INFORMATION (continued on next page)

Contraindications

EUCRISA is contraindicated in patients with known hypersensitivity to crisaborole or any component of the formulation.

Warnings and Precautions

Hypersensitivity reactions, including contact urticaria, have occurred in patients treated with EUCRISA and should be suspected in the event of severe pruritus, swelling, and erythema at the application site or at a distant site. Discontinue EUCRISA immediately and initiate appropriate therapy if signs and symptoms of hypersensitivity occur.

Please see Important Safety Information and Indication continued on next page.

Please see full Prescribing Information on the following pages or visit [EucrisaHCP.com](https://www.eucrisahcp.com/).

<https://www.eucrisahcp.com/>

EUCRISA: THE ONLY FDA-APPROVED NONSTEROIDAL TOPICAL Rx AD MEDICATION THAT HAS ESTABLISHED SAFETY AND EFFECTIVENESS IN PATIENTS AS YOUNG AS 3 MONTHS¹

28-day, open-label, single-arm study designed to evaluate the safety of EUCRISA in infants (N=137)

Safety outcomes with EUCRISA in patients aged 3 to <24 months²

MOST FREQUENTLY REPORTED TREATMENT-RELATED TREATMENT AREA ADVERSE EVENTS (AEs) (≥2.5% of patients) ²		
Application Site Pain	Application Site Discomfort	Erythema*
3.6% (n=5)	2.9% (n=4)	2.9% (n=4)

- No new safety issues of concern were identified by laboratory findings, ECG, height, weight, or vital signs²

*Application site erythema was also reported in 2.2% (n=3) of patients.

Treatment emergent adverse events (TEAE)

- The most frequently reported (≥5%) all-cause TEAEs were pyrexia (9.5%), upper respiratory tract infection (7.3%), diarrhea (7.3%), dermatitis atopic (6.6%), dermatitis diaper (6.6%), and cough (5.1%)²
- Discontinuation rate due to TEAEs was 2.9% (n=4) for EUCRISA (all remained in the study)²
 - One patient experienced a serious TEAE of febrile convulsion (not related to treatment)²

28-day infant open-label, single-arm safety study in which efficacy was an exploratory endpoint (N=137)

Efficacy outcomes with EUCRISA in patients aged 3 to <24 months²

- 47.3% of patients had an ISGA Clear (0) or Almost Clear (1) at day 29
- 57.5% decrease in mean EASI score from baseline to day 29

ISGA=Investigator's Static Global Assessment; EASI=Eczema Area and Severity Index.

Study Design & Limitations²

- Use of EUCRISA in this age group is supported by evidence from one multicenter, open-label, single-arm 28-day trial, which included 137 infant patients aged 3 months to <24 months
- Outcomes were reported by a caregiver or investigator



Prescribe EUCRISA for your appropriate mild-to-moderate AD patients as young as 3 months of age

IMPORTANT SAFETY INFORMATION (continued)

Adverse Reactions

The most common treatment-related adverse reaction occurring in clinical trials was application site pain, such as burning or stinging.

INDICATION

EUCRISA is indicated for topical treatment of mild-to-moderate atopic dermatitis in adult and pediatric patients 3 months of age and older.

Please see full Prescribing Information on the following pages or visit EucrisaHCP.com.

References: 1. EUCRISA® (crisaborole) Full Prescribing Information. April 2020. 2. Schlessinger J, Shepard JS, Gower R, et al. Safety, effectiveness, and pharmacokinetics of crisaborole in infants aged 3 to <24 months with mild-to-moderate atopic dermatitis: a phase IV open-label study (CrisADe CARE 1). *Am J Clin Dermatol*. 2020. <https://doi.org/10.1007/s40257-020-00510-6>. 3. Eichenfield LF, Cart RS, Forsha OW, et al. Long-term safety of crisaborole ointment 2% in children and adults with mild to moderate atopic dermatitis. *J Am Acad Dermatol*. 2017;77(4):641-649.eS. 4. Paller AS, Tom WL, Lebwohl MG, et al. Efficacy and safety of crisaborole ointment, a novel, nonsteroidal phosphodiesterase 4 (PDE4) inhibitor for the topical treatment of atopic dermatitis (AD) in children and adults. *J Am Acad Dermatol*. 2016;75(3):494-503.e 5. Data on file. Pfizer Inc., New York, NY.



HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use EUCRISA safely and effectively. See full prescribing information for EUCRISA.

EUCRISA® (crisaborole) ointment, for topical use
Initial U.S. Approval: 2016

RECENT MAJOR CHANGES

Indications and Usage (1) 03/2020

INDICATIONS AND USAGE

EUCRISA is a phosphodiesterase 4 inhibitor indicated for topical treatment of mild to moderate atopic dermatitis in adult and pediatric patients 3 months of age and older. (1)

DOSAGE AND ADMINISTRATION

- Apply a thin layer twice daily to affected areas. (2)
- For topical use only. (2)
- Not for ophthalmic, oral, or intravaginal use. (2)

DOSAGE FORMS AND STRENGTHS

Ointment, 2%. (3)

CONTRAINDICATIONS

Known hypersensitivity to crisaborole or any component of the formulation. (4)

WARNINGS AND PRECAUTIONS

Hypersensitivity reactions: If signs and symptoms of hypersensitivity occur, discontinue EUCRISA immediately and initiate appropriate therapy. (5.1)

ADVERSE REACTIONS

The most common adverse reaction occurring in $\geq 1\%$ in subjects is application site pain. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Pfizer Inc. at 1-800-438-1985 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.

Revised: 04/2020

FULL PRESCRIBING INFORMATION: CONTENTS*

- 1 INDICATIONS AND USAGE
- 2 DOSAGE AND ADMINISTRATION
- 3 DOSAGE FORMS AND STRENGTHS
- 4 CONTRAINDICATIONS
- 5 WARNINGS AND PRECAUTIONS
 - 5.1 Hypersensitivity Reactions
- 6 ADVERSE REACTIONS
 - 6.1 Clinical Trials Experience
 - 6.2 Postmarketing Experience
- 8 USE IN SPECIFIC POPULATIONS
 - 8.1 Pregnancy
 - 8.2 Lactation
 - 8.4 Pediatric Use
 - 8.5 Geriatric Use
- 11 DESCRIPTION

- 12 CLINICAL PHARMACOLOGY
 - 12.1 Mechanism of Action
 - 12.2 Pharmacodynamics
 - 12.3 Pharmacokinetics
- 13 NONCLINICAL TOXICOLOGY
 - 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
- 14 CLINICAL STUDIES
- 16 HOW SUPPLIED/STORAGE AND HANDLING
 - 16.1 How Supplied
 - 16.2 Storage and Handling
- 17 PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

EUCRISA is indicated for topical treatment of mild to moderate atopic dermatitis in adult and pediatric patients 3 months of age and older.

2 DOSAGE AND ADMINISTRATION

Apply a thin layer of EUCRISA twice daily to affected areas.

EUCRISA is for topical use only and not for ophthalmic, oral, or intravaginal use.

3 DOSAGE FORMS AND STRENGTHS

Ointment: 20 mg of crisaborole per gram (2%) of white to off-white ointment.

4 CONTRAINDICATIONS

EUCRISA is contraindicated in patients with known hypersensitivity to crisaborole or any component of the formulation. [see Warnings and Precautions (5.1)]

5 WARNINGS AND PRECAUTIONS

5.1 Hypersensitivity Reactions

Hypersensitivity reactions, including contact urticaria, have occurred in patients treated with EUCRISA.

Hypersensitivity should be suspected in the event of severe pruritus, swelling and erythema at the application site or at a distant site. If signs and symptoms of hypersensitivity occur, discontinue EUCRISA immediately and initiate appropriate therapy.

6 ADVERSE REACTIONS

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

In two double-blind, vehicle-controlled clinical trials (Trial 1 and Trial 2), 1012 subjects 2 to 79 years of age with mild to moderate atopic dermatitis were treated with EUCRISA twice daily for 4 weeks. The adverse reaction reported by $\geq 1\%$ of EUCRISA-treated subjects is listed in Table 1.

Table 1: Adverse Reaction Occurring in $\geq 1\%$ of Subjects in Atopic Dermatitis Trials through Week 4

Adverse Reaction	EUCRISA N=1012 n (%)	Vehicle N=499 n (%)
Application site pain ^a	45 (4)	6 (1)

^a Refers to skin sensations such as burning or stinging.

Less common (<1%) adverse reactions in subjects treated with EUCRISA included contact urticaria [see Warnings and Precautions (5.1)].

6.2 Postmarketing Experience

The following adverse reactions have been identified during postapproval use of EUCRISA. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure:

Skin and Subcutaneous: allergic contact dermatitis

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

There is no available data with EUCRISA in pregnant women to inform the drug-associated risk for major birth defects and miscarriage. In animal reproduction studies, there were no adverse developmental effects observed with oral administration of crisaborole in pregnant rats and rabbits during organogenesis at doses up to 3 and 2 times, respectively, the maximum recommended human dose (MRHD) [*see Data*].

The background risk of major birth defects and miscarriage for the indicated population is unknown. All pregnancies carry some risk of birth defect, loss, or other adverse outcomes. The background risk of major birth defects in the U.S. general population is 2% to 4% and of miscarriage is 15% to 20% of clinically recognized pregnancies.

Data

Animal Data

Rat and rabbit embryo-fetal development was assessed after oral administration of crisaborole. Crisaborole did not cause adverse effects to the fetus at oral doses up to 300 mg/kg/day in pregnant rats during the period of organogenesis (3 times the MRHD on an area under the curve (AUC) comparison basis). No crisaborole-related fetal malformations were noted after oral treatment with crisaborole in pregnant rats at doses up to 600 mg/kg/day (13 times the MRHD on an AUC comparison basis) during the period of organogenesis. Maternal toxicity was produced at this high dose of 600 mg/kg/day in pregnant rats and was associated with decreased fetal body weight and delayed skeletal ossification. Crisaborole did not cause adverse effects to the fetus at oral doses up to the highest dose tested of 100 mg/kg/day in pregnant rabbits during the period of organogenesis (2 times the MRHD on an AUC comparison basis).

In a prenatal/postnatal development study, pregnant rats were treated with crisaborole at doses of 150, 300, or 600 mg/kg/day by oral gavage during gestation and lactation (from gestation day 7 through day 20 of lactation). Crisaborole did not have any adverse effects on fetal development at doses up to 300 mg/kg/day (3 times the MRHD on an AUC comparison basis). Maternal toxicity was produced at the high dose of 600 mg/kg/day in pregnant rats and was associated with stillbirths, pup mortality, and reduced pup weights.

8.2 Lactation

Risk Summary

There is no information available on the presence of EUCRISA in human milk, the effects of the drug on the breastfed infant or the effects of the drug on milk production after topical application of EUCRISA to women who are breastfeeding. EUCRISA is systemically absorbed. The lack of clinical data during lactation precludes a clear determination of the risk of EUCRISA to a breastfed infant. Therefore, the developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for EUCRISA and any potential adverse effects on the breastfed infant from EUCRISA or from the underlying maternal condition.

8.4 Pediatric Use

The safety and effectiveness of EUCRISA have been established in pediatric patients ages 3 months and older for topical treatment of mild to moderate atopic dermatitis. Use of EUCRISA in this age group is supported by data from two 28-day adequate, vehicle-controlled safety and efficacy trials which included 1,313 pediatric subjects ages 2 years to 17 years of whom 874 received EUCRISA. The most commonly reported adverse reaction in subjects 2 years and older was application site pain. Additionally, use of EUCRISA in pediatric patients ages 3 months to less than 2 years was supported by data from a 28-day open-label, safety and pharmacokinetics (PK) trial in 137 subjects. No new safety signals were identified in subjects 3 months to less than 2 years of age [see *Adverse Reactions (6.1)*, *Clinical Pharmacology (12.3)*, and *Clinical Studies (14)*].

The safety and effectiveness of EUCRISA in pediatric patients below the age of 3 months have not been established.

8.5 Geriatric Use


Clinical studies of EUCRISA did not include sufficient numbers of subjects age 65 and over to determine whether they respond differently from younger subjects.

11 DESCRIPTION

EUCRISA contains 2% crisaborole (w/w) in a petrolatum-based, white to off-white ointment and is for topical use. The active ingredient, crisaborole, is a phosphodiesterase-4 (PDE-4) inhibitor.

Crisaborole is described chemically as 5-(4-cyanophenoxy)-1,3-dihydro-1-hydroxy-[2,1]-benzoxaborole. The empirical formula is $C_{14}H_{10}BNO_3$ and the molecular weight is 251.1 g/mol.

The structural formula is represented below:



Crisaborole drug substance is freely soluble in common organic solvents such as isopropyl alcohol and propylene glycol, and insoluble in water.

Each gram of EUCRISA contains 20 mg of crisaborole in an ointment containing white petrolatum, propylene glycol, mono- and di-glycerides, paraffin, butylated hydroxytoluene, and edetate calcium disodium.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Crisaborole is a phosphodiesterase 4 (PDE-4) inhibitor. PDE-4 inhibition results in increased intracellular cyclic adenosine monophosphate (cAMP) levels. The specific mechanism(s) by which crisaborole exerts its therapeutic action for the treatment of atopic dermatitis is not well defined.

12.2 Pharmacodynamics

Cardiac Electrophysiology

At therapeutic doses, EUCRISA ointment is not expected to prolong QTc to any clinically relevant extent.

12.3 Pharmacokinetics

Absorption

The PK of EUCRISA were investigated in 33 pediatric subjects 2 to 17 years of age with mild to moderate atopic dermatitis and a mean \pm SD body surface area (BSA) involvement of $49 \pm 20\%$ (range 27% to 92%). In this study, subjects applied approximately 3 mg/cm² of EUCRISA ointment (dose range was approximately 6 g to 30 g per application) twice daily for 8 days.

Plasma concentrations were quantifiable in all the subjects. The mean \pm SD maximum plasma concentration (C_{\max}) and area under the concentration time curve from 0 to 12 hours post dose (AUC_{0-12}) for crisaborole on Day 8 were 127 ± 196 ng/mL and 949 ± 1240 ng h/mL, respectively. Systemic concentrations of crisaborole were at steady state by Day 8. Based on the ratios of AUC_{0-12} between Day 8 and Day 1, the mean accumulation factor for crisaborole was 1.9.

The PK of EUCRISA were investigated in 13 subjects 4 months to less than 24 months of age. The mean \pm SD C_{\max} and AUC_{0-12} for crisaborole were 188 ± 100 ng/mL and 1164 ± 550 ng·h/mL, respectively.

Distribution

Based on an in vitro study, crisaborole is 97% bound to human plasma proteins.

Elimination

Metabolism

Crisaborole is substantially metabolized into inactive metabolites. The major metabolite 5-(4-cyanophenoxy)-2-hydroxyl benzylalcohol (metabolite 1), is formed via hydrolysis; this metabolite is further metabolized into downstream metabolites, among which 5-(4-cyanophenoxy)-2-hydroxyl benzoic acid (metabolite 2), formed via oxidation, is also a major metabolite.

PK of metabolites 1 and 2 were assessed in the PK study described above and the systemic concentrations were at or near steady state by Day 8. Based on the ratios of AUC_{0-12} between Day 8 and Day 1, the mean accumulation factors for metabolites 1 and 2 were 1.7 and 6.3, respectively.

Excretion

Renal excretion of metabolites is the major route of elimination.

Drug Interaction Studies

In vitro studies using human liver microsomes indicated that under the conditions of clinical use, crisaborole and metabolite 1 are not expected to inhibit cytochrome P450 (CYP) 1A2, 2B6, 2C8, 2C9, 2C19, 2D6, and 3A4.

In vitro human liver microsomes studies for metabolite 2 showed that it did not inhibit activities of CYP2C19, 2D6, and 3A4; was a weak inhibitor of CYP1A2 and 2B6; and a moderate inhibitor of CYP2C8 and 2C9. The most sensitive enzyme, CYP2C9, was further investigated in a clinical trial using warfarin as a CYP2C9 substrate. The results of this study showed no drug interaction potential.

In vitro studies in human hepatocytes showed that under the conditions of clinical use, crisaborole and metabolites 1 and 2 are not expected to induce CYP enzymes.

In vitro studies showed that crisaborole and metabolite 1 did not inhibit the activities of uridine diphosphate (UDP)-glucuronosyltransferase (UGT) 1A1, 1A4, 1A6, 1A9, 2B7, and 2B15. Metabolite 2 did not inhibit UGT1A4, 1A6, 2B7, and 2B15. Metabolite 2 showed weak inhibition of UGT1A1, however, no clinically

significant drug interactions are expected between crisaborole (and its metabolites) and UGT1A1 substrates at therapeutic concentrations. Metabolite 2 showed moderate inhibition of UGT1A9 and may result in a moderate increase of the concentrations of sensitive UGT1A9 substrates.

In vitro studies indicate that under the condition of clinical use, crisaborole and metabolites 1 and 2 are not expected to cause clinically significant interactions with substrates of P-glycoprotein and organic anionic or cationic transporters. Crisaborole and metabolite 1 are not expected to inhibit breast cancer resistance protein (BCRP); metabolite 2 is expected to inhibit BCRP at therapeutic concentrations.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

In an oral carcinogenicity study in Sprague-Dawley rats, oral doses of 30, 100, or 300 mg/kg/day crisaborole were administered to rats once daily. A crisaborole-related increased incidence of benign granular cell tumors in the uterus with cervix and vagina (combined) was noted in 300 mg/kg/day crisaborole treated female rats (2 times the MRHD on an AUC comparison basis). The clinical relevance of this finding is unknown.

In a dermal carcinogenicity study in CD-1 mice, topical doses of 2%, 5%, or 7% crisaborole ointment were administered once daily. No crisaborole-related neoplastic findings were noted at topical doses up to 7% crisaborole ointment (1 times the MRHD on an AUC comparison basis).

Crisaborole revealed no evidence of mutagenic or clastogenic potential based on the results of two in vitro genotoxicity tests (Ames assay and human lymphocyte chromosomal aberration assay) and one in vivo genotoxicity test (rat micronucleus assay).

No effects on fertility were observed in male or female rats that were administered oral doses up to 600 mg/kg/day crisaborole (13 times the MRHD on an AUC comparison basis) prior to and during early pregnancy.

14 CLINICAL STUDIES

Two multicenter, randomized, double-blind, parallel-group, vehicle-controlled trials (Trials 1 and 2) treated a total of 1522 subjects 2 to 79 years of age (86.3% of subjects were 2 to 17 years of age) with a 5% to 95% treatable BSA. At baseline, 38.5% of the subjects had an Investigator's Static Global Assessment [ISGA] score of 2 (mild), and 61.5% had an ISGA score of 3 (moderate), in the overall assessment of atopic dermatitis (erythema, induration/papulation, and oozing/crusting) on a severity scale of 0 to 4.

In both trials, subjects were randomized 2:1 to receive EUCRISA or vehicle applied twice daily for 28 days. The primary efficacy endpoint was the proportion of subjects at Day 29 who achieved success, defined as an ISGA grade of Clear (score of 0) or Almost Clear (score of 1) with a 2-grade or greater improvement from baseline, comparing EUCRISA-treated subjects to vehicle-treated subjects.

Efficacy results from the two trials are summarized in Table 2.


Table 2: Primary Efficacy Outcomes in Subjects with Mild to Moderate Atopic Dermatitis at Day 29

	Trial 1		Trial 2	
	EUCRISA (N=503)	Vehicle (N=256)	EUCRISA (N=513)	Vehicle (N=250)
Success in ISGA^a	32.8%	25.4%	31.4%	18.0%

^a Defined as an ISGA score of Clear (0) or Almost Clear (1) with a 2-grade or greater improvement from baseline.

The success rates over time are presented in Figure 1.

Figure 1: Success in ISGA^a Over Time in Subjects with Mild to Moderate Atopic Dermatitis



^a Success is defined as an ISGA score of Clear (0) or Almost Clear (1) with a 2-grade or greater improvement from baseline.

16 HOW SUPPLIED/STORAGE AND HANDLING

16.1 How Supplied

EUCRISA is a white to off-white ointment containing 2% crisaborole and is supplied in 60 g and 100 g laminate tubes.

60 g tube: NDC 55724-211-21

100 g tube: NDC 55724-211-11

16.2 Storage and Handling

Store at 20°C–25°C (68°F–77°F); excursions permitted to 15°C–30°C (59°F–86°F).
[see USP Controlled Room Temperature].

Keep tube tightly closed.

17 PATIENT COUNSELING INFORMATION

Advise the patient or caregivers to read the FDA-approved patient labeling (Patient Information).

Hypersensitivity Reactions

Advise patients to discontinue EUCRISA and seek medical attention immediately if signs or symptoms of hypersensitivity occur [see Warnings and Precautions (5.1)].

Administration Instructions

Advise patients or caregivers that EUCRISA is for external use only and is not for ophthalmic, oral, or intravaginal use.

This product's labeling may have been updated. For the most recent prescribing information, please visit www.pfizer.com.



Distributed by

Pfizer Labs

Division of Pfizer Inc, NY, NY 10017

LAB-0916-5.0

PATIENT INFORMATION
EUCRISA® (you-KRIS-a)
(crisaborole) ointment, 2%

Important information: EUCRISA is for use on skin (topical use) only. Do not use EUCRISA in your eyes, mouth, or vagina.

What is EUCRISA?

EUCRISA is a prescription medicine used on the skin (topical) to treat mild to moderate eczema (atopic dermatitis) in adults and children 3 months of age and older.

It is not known if EUCRISA is safe and effective in children under 3 months of age.

Who should not use EUCRISA?

Do not use EUCRISA if you are allergic to crisaborole or any of the ingredients in EUCRISA. See the end of this leaflet for a complete list of ingredients in EUCRISA.

Before using EUCRISA, tell your healthcare provider about all of your medical conditions, including if you:

- are pregnant or plan to become pregnant. It is not known if EUCRISA will harm your unborn baby.
- are breastfeeding or plan to breastfeed. It is not known if EUCRISA passes into your breast milk.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

How should I use EUCRISA?

- Use EUCRISA exactly as your healthcare provider tells you to use it.
- Apply a thin layer of EUCRISA to the affected areas 2 times each day.
- Wash your hands after applying EUCRISA, unless hands are being treated. If someone else applies EUCRISA for you, they should wash their hands after applying EUCRISA.

What are the possible side effects of EUCRISA?

EUCRISA may cause side effects.

- **Allergic reactions.** EUCRISA may cause allergic reactions at or near the application site. These can be serious and may include hives, itching, swelling, and redness. If you have any of these symptoms, stop using EUCRISA and get medical help right away.

The most common side effect of EUCRISA is application site pain, such as burning or stinging.

This is not the only possible side effect of EUCRISA.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store EUCRISA?

- Store EUCRISA at room temperature, between 68°F and 77°F (20°C and 25°C).
- Keep the tube tightly closed.

General information about the safe and effective use of EUCRISA

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use EUCRISA for a condition for which it was not prescribed. Do not give EUCRISA to other people, even if they have the same symptoms that you have. It may harm them. You can ask your pharmacist or healthcare provider for information about EUCRISA that is written for healthcare professionals.

What are the ingredients in EUCRISA?

Active ingredient: crisaborole

Inactive ingredients: white petrolatum, propylene glycol, mono- and di-glycerides, paraffin, butylated hydroxytoluene, and edetate calcium disodium.

For more information, call 1-866-EUCRISA [1-866-382-7472] or go to www.EUCRISA.com



LAB-0917-4.0

This Patient Information has been approved by the U.S. Food and Drug Administration.

Revised: 04/2020